


Finding YSOs in GLIMPSE

Thomas Robitaille


Barb Whitney, Remy Indebetouw, Kenneth Wood, Pia Denzmore



University
of
St Andrews



Low-mass star formation paradigm



Class 0

$$t \sim 10^4 \text{ yr}$$

$$R_{\text{env}} \sim 10^4 \text{ AU}$$

Class I

$$t \sim 10^5 \text{ yr}$$

$$R_{\text{env}} \sim 10^3 \text{ AU}$$

Class II


$$t \sim 10^6 \text{ yr}$$

$$R_{\text{disk}} \sim 100 \text{ AU}$$


Class III

$$t \sim 10^7 \text{ yr}$$


Example of a YSO SED



Example of a YSO SED



Example of a YSO SED





IRAC bands [4] (R), [2] (G), [3] (B)



MIPS 24 μ (R), IRAC bands [4] (G), [I] (B)


Predicting colors of YSOs

- Use radiation transfer models to predict how YSOs at various evolutionary stages should look.
- We have done this by producing synthetic SEDs for 20,000 different combinations of physical parameters and 10 inclinations
- These are available online


The new set of models

- Use Monte-Carlo radiation transfer models (Whitney et al.)
- Span a wide range of stellar masses & evolutionary states
- Allow for uncertainties in evolutionary stages by allowing ranges of parameters
- Sample randomly within these ranges



The general YSO model













Sampling the model parameters...




Sampling the model parameters...








200,000 models at a glance...




1000 SEDs

200,000 models at a glance...



100,000 SEDs



200,000 models at a glance...



200,000 SEDs

Spectral indices

- Easiest way to quantify observations of a YSO: define a *spectral index* between two wavelengths




Spectral indices

- Defined as $\alpha = \frac{d \log \lambda F_\lambda}{d \log \lambda}$
- Traditionally three 'Classes' of objects:


$\alpha > 0$	$\longrightarrow$	Class I
$-2 < \alpha < 0$	$\longrightarrow$	Class II
$-3 < \alpha < -2$	$\longrightarrow$	Class III

Spectral index vs YSO evolution?

$$\alpha = \frac{d \log \lambda F_{\lambda}}{d \log \lambda}$$



Spectral indices (predictions)




IRAC 3.6 to 8.0 μm

J to IRAC 8.0 μm

K to MIPS 24 μm

Spectral indices (observations)



↑ ↑
YSOs

Stars

↑ ↑
YSOs


Stars

Lada et al. (2006)

Spectral index vs YSO evolution?



- 2-25 μ m spectral index is a decent (although not perfect) indicator of evolutionary stage.
- However, due to the small wavelength range, IRAC does not always correlate with evolutionary stage.

IRAC vs 2-25 μ m




Color-color plots

- Use three or four datapoints to define *colors*




- Find magnitude at each of these points

YSOs in IRAC+MIPS color-color plots



Robitaille et al. (2006)

Available YSO model SEDs



Summary

- Can use spectral indices or color-color diagrams to find YSOs in GLIMPSE.
- MIPS 24 μ m is useful (even upper limits on fluxes).
- Synthetic SEDs for a large range of physical parameters are available online:

<http://www.astro.wisc.edu/protostars>